

Research Article

Influence of zeatin, glutamin and auxins on root and shoot organogenesis of Guava (*Psidium guajava* L.) cv. *safeda* seedling explants

Roshan Zamir¹, Abdur Rab², Muhamad Sajid² and Izhar Ahmad^{3*}

1. Nuclear Institute for Food & Agriculture (NIFA), Peshawar-Pakistan

2. Department of Horticulture, The University of Agriculture, Peshawar-Pakistan

3. Department of Botany, Islamia College Peshawar-Pakistan

*Corresponding author's email: izhar.ahmad@icp.edu.pk

Citation

Roshan Zamir, Abdur Rab, Muhamad Sajid and Izhar Ahmad. Influence of zeatin, glutamin and auxins on root and shoot organogenesis of Guava (*Psidium guajava* L.) cv. *safeda* seedling explants. Pure and Applied Biology. Vol. 6, Issue 1, pp197-206. <http://dx.doi.org/10.19045/bspab.2017.60014>

Received: 07/01/2017

Revised: 02/02/2017

Accepted: 05/02/2017

Online First: 07/02/2017

Abstract

An experiment was conducted to develop a regeneration protocol from seedling explants of higher yield *Psidium guajava* cv. *safeda*. Initially surface sterilized guava seeds were germinated inside the conical flasks. Explants taken from 49-day old seedlings were cultured on MS-medium modified with different concentration of plant growth regulators. Maximum explants showing shoot response (48%) was observed on MS-medium supplemented with 6-benzyladenin (BA) 1.0 mg l⁻¹ combined with 0.5 mg l⁻¹ zeatin. On the same media after 38-days of culturing, maximum number of 3.6 shoots per explant, 2.6 cm long was also recorded. On MS containing combination of BA (2.0 mg l⁻¹) and zeatin (1.0 mg l⁻¹), 37% explants responded which produced 2.3 shoot per explant after 42 days of culturing. Whereas on MS containing glutamin 250 mg l⁻¹, 25% explants responded and gave 2.2 shoots per explants after 46-days. However, when the concentration of glutamin alone was increased from 250 to 500 mg l⁻¹ the shoot formation potential of explant decreased to 10%. Addition of BA, glutamin and zeatin alone to MS medium was less effective. However, combination of BA with zeatin was effective in shooting response, number of shoot/explants and shoot length. Maximum number of explants (76.6%) responded to rooting with 1.5 mg l⁻¹ IAA in combination with 1.0 mg l⁻¹ NAA. On average, maximum of (2.5) roots were also recorded per explants in 42 days. *In vitro* plantlets were acclimatized and shifted to soil successfully.

Keywords: *Psidium guajava* L; Micropropagation; Zeatin; Glutamin and auxins

Introduction

Guava (*Psidium guajava* L.) is one of the most common fruit crops of Pakistan which ranks third after citrus and mango in area and production wise [1]. Guava was originated in tropical America stretching

from Mexico to Peru [2]. Guava is grown throughout the tropical and subtropical regions of the world but the average yield, however, is much less than its potential, probably because the orchards established from seedling plants [3]. The importance of

guava is increasing due to its nutritional value, biannual bearing and affordable prices [4]. It is a well-known fruit of tropical and subtropical regions, where it is consumed fresh or processed into different products [5]. Guava is usually propagated by seeds and natural cross pollination is responsible for diversity in seedlings plants [4]. Guava propagation by layering, inarching, and budding is achievable but cumbersome and time consuming [6]. During the past three decades, several techniques have been practiced for propagation of guava but no one could be perfected. Guava is recalcitrant specie for tissue culture when the explants are taken from mature part of the plant [7]. It is difficult to establish aseptic cultures from field grown mature woody plants due to endophytic microorganisms [8].

Clonal propagation of guava through micropropagation has been demonstrated using seedling explants, shoot tips and nodal segments [9-11]. Micropropagation of guava has been achieved by single node cuttings from seedlings raised in green house and *in vitro* raised seedlings [12]. A regeneration protocol from hypocotyl seedling explants of guava was also carried out by [9]. Micropropagation can be exploited to produce guaranteed quality, reliable and true to type plants for farmer community. It is rapid, efficient and cost effective technology, which will help in germplasm preservation and genetic improvement of the crop. Moreover, true to type plants can be produced throughout the year for the growers through micropropagation. The present studies were carried out to investigate rapid production of true-to-type plants *in vitro* from seedling explants of elite *Guava cv. safeda*.

Materials and methods

Seed extraction and sterilization

Seeds from well ripen fruits of *Guava cv. safeda* were extracted. After thoroughly

washing, seeds were surface sterilized with 0.1% Mercuric Chloride for 5 minutes followed by four rinsing with autoclaved sterile distilled water according to [4]. These were cultured in 500 ml conical flasks inside laminar flow bench on sterilized filter paper laid on cotton below. The flasks were kept in the dark for the first three weeks and then shifted to 16 hour photoperiod produced by white fluorescent lamps. Forty-nine-day old contamination free *in vitro* seedlings were used as explants source.

Explants sterilization, culturing and media composition

The explants (0.5 - 1cm) were aseptically excised from the seedlings and given a quick rinse with 70% ethanol followed by two rinses with sterile distilled water. The excised explants were cultured according to [13] containing 6- benzyladenine (BA) (1.0 and 2.0 mg l⁻¹) and zeatin (0.5 and 1.0 mg l⁻¹) alone and in combinations. One combination of MS media was also supplemented with glutamin 250 and 500 mg l⁻¹. Solidification of media was done with 0.8% agar. The media was also supplemented with 30 g l⁻¹ sucrose. The pH of the medium was adjusted with pH meter (Inolab Germany) to 5.80 before autoclaving at (1.02 kg/cm² for 15 min). Cultures were kept in dark on (25 ± 2 °C) for initial 24 hours and then shifted to light from cool white fluorescent lamps (3000 Lux) maintained at 16 hours photoperiod.

Shoot organogenesis

The cultures were observed for regeneration of shoot and shoot multiplication. The parameters on days to shooting, % shooting, number of shoot/explants and length of microshoot were recorded on regeneration media after 8-9 weeks of culturing.

Root organogenesis

The Micro shoots produced from seedling explants on MS media supplemented with different cytokinin were excised and shifted to rooting medium containing different

concentrations and combinations of Indol-3-butyric acid (IBA), Indole-3-acetic acid (IAA) and Naphthalene acetic acid (NAA). The parameters on % rooting, days to rooting, number of roots/shoot and root length was recorded after 7-8 weeks of culturing.

Acclimatization of plantlets

When the micro shoot rooted successfully they were thoroughly washed with sterile doubled distilled water and transplanted in to small pots containing mixture of sterile soil of sand, silt and clay at (1:1:1). They were covered with transparent polythene sheet to arrest humidity. The hardened plantlets (35-45 days) with 2-4 fully expanded leaves were again shifted to 7.5cm diameter plastic pots containing sand, silt and clay. They were then transferred to the glasshouse. The plants were covered with polyethylene bags for 15-20 days. These pots were kept at 25°C in 16 hour photoperiod for 3-4 weeks before transferring to lathe house.

Experimental design

The experiment was laid out in Completely Randomized Design with three replications. The data were statistically analyzed by using the software Stat. 8.1 USA analysis of variance. Duncan's Multiple Range test was used to compare the treatments among themselves.

Results and discussion

Percent shooting

The data presented in figure-1 revealed that different PGRs have significantly affected percent shootings in seedling explants of guava. Explants cultured on MS with 1.0 mg^l⁻¹ BA in combination with 0.5 mg^l⁻¹ zeatin gave maximum (48%) shooting. With the same combinations when the concentration of cytokinin was doubled the shooting frequency reduced to 37%. Twenty two percent explants on MS medium with 1.0 mg^l⁻¹ BA responded to shoot regeneration after seven weeks of culturing.

However, an increase in BA from 1.0 mg^l⁻¹ to 2.0 mg^l⁻¹ decreased shoot regeneration potential of explants. Compared to 22% explants regeneration, only 15% explants responded. Only 25% explants responded to shoot regeneration when glutamin 250 mg^l⁻¹ was added to MS medium. All concentration of zeatin and BA alone gave less shooting frequency as compared to their combinations. On the other hand, lower concentration of glutamin (250 mg^l⁻¹) gave higher response in shoot regeneration as compared to the higher concentration (500 mg^l⁻¹) which gave only 10% shooting response (Fig.1). When cytokinins alone were used, the shooting frequency of explants decreases. However, combination of BA with zeatin was effective in shooting response, number of shoot/explants and shoot length.

Most of the literature on micro propagation indicates that cytokinin and other hormone play a key role in shoot organogenesis in guava [14-16]. The present study reveals that combination of cytokinins had encouraged regeneration and it was almost doubled as compared to their use alone. Lower concentrations of BA combined with zeatin have proved to be the best one in the regeneration and shoot induction of guava. In contrast to [10], maximum number of shoot per explant in seedling tip of guava was reported when the medium was supplemented with 2.0 mg^l⁻¹ BA. Similar results were recorded by [9, 11] of seedling explants regeneration on Ms-medium supplemented with lower concentration of BA (0.01-1.0 mg^l⁻¹). Many researchers suggested low level of cytokinins in regeneration of guava for proper growth of leaves and uniform growth in micro-shoots [11, 17, 18]. It was reported that prolonged exposure to high level of cytokinin caused necrosis in leaves [6, 11]. BA is a powerful cytokinin in shooting of guava and its use has been reported by several researchers [11,

17, 19]. glutamin and tryptophan had no positive effect on the tissue culture system of upland rice cultivars (except cultivar Lamsan) regarding callus induction [20].

These results strongly support the findings of our results that higher concentration of glutamin gave lower response (10%) only.




Figure 1. Effect of PGRs on % shooting. Means with same letters are not significantly different at $P < 0.05$ using DMR test

Days to shoot induction

The data presented in figure-2 revealed that the lowest number of days (39) was recorded when MS media was augmented with 1.0 mg l^{-1} BA combined with 0.5 mg l^{-1} zeatin followed by 41.5 days on the medium containing 2.0 mg l^{-1} BA combined with 1.0 mg l^{-1} zeatin. The highest number of days (51.8) to shooting was recorded when MS medium was supplemented with 1.0 mg l^{-1} zeatin. It followed with 51 days on media with 2.0 mg l^{-1} BA and 50 days on media with 500 mg l^{-1} glutamin. The present studies reveal that combination of BA and zeatin has encouraged early shooting. Higher concentrations of BA and zeatin alone have delayed shoot induction. In earlier studies on guava shoot tips and nodal explants, it was recorded 4-6 weeks for shoot induction on MS medium supplemented with $4.5 \mu\text{M}$ BA [21]. In contrast, [11] recorded the

maximum number of days (52.5) and the minimum days to shoot induction (35.8) in seedling explant of guava on $2.22 \mu\text{M}$ BAP and TDZ + NAA. The earliness in shooting in this study might be due to different media formulation (incorporation of NAA and TDZ) or could be due to explants difference. However, single shoot formation per explants was recorded on medium containing BA at $0.01\text{-}1 \text{ mg l}^{-1}$ after 8 weeks of culturing and higher concentrations of cytokinins have created negative effects on the regeneration of guava *in vitro* [17]. In case of our studies the combination of BA with zeatin in both concentrations has reduced the number of days to shoot induction. It is likely due the fact that both cytokinin play a key role in regeneration and their combined effect have synergistically reduced the number of days to shoot regeneration.




Figure 2. Effect of PGRs on days to shoot formation. Means with same letters are not significantly different at $P < 0.05$ using DMR test

Number of shoot per explant

Maximum average number of shoots (3.6) per explants was recorded on medium supplemented with 1.0 mg l^{-1} BA in combination with 0.5 mg l^{-1} zeatin (Fig-3). This was followed by BA alone at 1.0 mg l^{-1} with 3.3 shoots per explants. This parameter once again confirms the effectiveness of BA irrespective of its combination with other cytokinin. The findings of our results is confirmed from the earlier studies of [9, 10, 19] who reported that multiple shoots were induced to form by enhancement of axillary branching with BA ($1.0\text{-}2.0 \text{ mg l}^{-1}$) without addition of any auxin and gibberellins. In contrast, [11] reported maximum of 5.2 shoot per explants when BAP and kinetin were incorporated in MS medium at 2.2 and $2.32 \mu\text{M}$ respectively. They further observed that BAP was effective alone in lower

concentrations for producing shoots. The high shooting frequency in this study may be due to incorporation of Kinitin or likely due to varietal difference. In our studies, glutamin at 500 mg l^{-1} and zeatin at both concentrations (0.5 and 1.0 mg l^{-1}) produced single shoot on average. However, the average number of shoot per explants was doubled when the concentration of glutamin was decreased from 500 to 250 mg l^{-1} . These results are in line with those reported [22] who investigated the effect of glutamin and asparagine on the growth and metabolic changes in French bean under *in vitro* conditions and observed that all growth parameters were increased by 1 and 2 mM glutamin or asparagine but when the concentration was increased the regeneration response was decreased.




Figure 3. Effect of PGRs on number of shoots/explant. Means with same letters are not significantly different at $P < 0.05$ using DMR test

Shoot length (cm)

The data in figure-4 revealed that on average, all concentrations of BA produced 1.5 cm long shoot in seven weeks. However, the longest shoot of 2.3 cm was produced when 1.0 mg l^{-1} BA was combined with 0.5 mg l^{-1} zeatin. This was followed by combination of BA (2.0 mg l^{-1}) and zeatin (1.0 mg l^{-1}). zeatin alone at both concentrations (0.5 and 1.0 mg l^{-1}) produced 0.8 and 1.0 cm long shoots. The most inferior results in case of shoot length was recorded when the medium was supplemented with glutamin 250 and 500 mg l^{-1} with maximum of 0.6 cm long shoot. The longest shoots were also recorded in

guava from green house grown explants by the application of lower concentrations (0.25 mg l^{-1}) of BAP [17] while the longest shoot of 2.5 cm on medium containing BAP $2.2 \mu\text{M}$ +kinitin $0.93 \mu\text{M}$ was achieved by [11]. In our studies all the concentrations of BA alone and in combination with zeatin was effective in shoot length. Among different cytokinin used for the proliferation of shoots from seedlings explants of guava, BA and zeatin was found superior to all. Similar results has been found during *in vitro* regeneration of many tropical fruit species such as guava [23], jack fruit [24-26], carambola [27], pomegranate [28] and jujube [29].




Figure 4. Effect of PGRs on average length of microshoot. Means with same letters are not significantly different at $P < 0.05$ using DMR test

Percent rooting

The percent rooting among all the treatments tried, maximum number of plantlets showing roots (76.6%) was observed on MS-medium containing 1.5 mg l^{-1} IAA and 1.0 mg l^{-1} NAA after 42 days. This was followed by 1.0 mg l^{-1} IAA and 0.5 mg l^{-1} NAA with 76% rooting. However, when the concentration of IBA increased to 1.5 mg l^{-1} with 1.0 mg l^{-1} NAA, reduced rooting to 73.3% of micro shoots. Again when the concentration of IBA decreased from 1.5 to 1.0 mg l^{-1} and NAA from 1.0 to 0.5 mg l^{-1} , rooting response decreased to 70%. Most of the relevant literature reported on the application of auxin indicates that IAA is very effective in various species of tropical trees [14], [30- 33]. In our studies at both concentrations of IAA or IBA, rooting response increases when NAA increase from 0.5 to 1.0 mg l^{-1} . These results are confirmed by those reported by [11, 32] on the rooting of guava *in vitro*. In contrast, the highest

rooting response of 95.9% on medium containing IBA $4.90 \mu\text{M}$ + 100 mg l^{-1} activated charcoal was reported by [11]. The high response of rooting in this case may be due the addition of activated charcoal in the medium. Addition of 100 mg l^{-1} activated charcoal had adsorbed toxins and phenolic compounds resulting in high rooting frequency.

Days to rooting

The data presented in table-1 revealed that among all the combinations, the lowest numbers of days (42) were taken to roots when the medium was supplemented with either 1.0 or 1.5 mg l^{-1} IAA or 0.5 or 1.0 mg l^{-1} NAA. It was followed by 1.5 mg l^{-1} IBA + 1.0 mg l^{-1} NAA with (45) days. The highest numbers of days (49) were taken to roots by the medium containing 1.0 mg l^{-1} IBA + 0.5 mg l^{-1} NAA. From present studies it is clear that IAA has played role and seems effective in reducing days to rooting. The lowest days of 22.5 to rooting in seedling explants

of guava on medium containing IBA 4.90 μM combined with activated charcoal 100 mg l^{-1} was reported by [11]. However, the highest number of days taken to rooting was 38.9 on the medium that containing 2.46 μM

IBA. The lowest days to rooting than our studies may be due to the addition of activated charcoal in the medium.

Table 1. Response of micro shoot from seedling explants of guava to rooting using various concentrations and combinations of auxin

Auxins (mg l^{-1})	Days to root induction	% root response	No. of roots/shoot
IBA 1.0 +NAA 0.5	49 a	70.0 c	1.9 d
IBA 1.5+NAA 1.0	45 b	73.3 b	2.2 b
IAA 1.0 + NAA0.5	42 c	76.0 a	2.0 c
IAA 1.5 + NAA1.0	42 c	76.6 a	2.5 a
LSD	2.2	1.2	0.15

Means not followed by same letters are significantly different at $P < 0.05$ using LSD test

Number of roots per shoot

Maximum number of roots per shoot (2.5) was recorded on medium containing 1.5 mg l^{-1} IAA + 1.0 mg l^{-1} NAA followed by 2.2 roots per shoot on the medium contained 1.5 mg l^{-1} IBA + 1.0 mg l^{-1} NAA (Table-1). Lower concentrations of NAA irrespective of combination with IBA or IAA have decreased the number of roots per explants. The number of roots per shoots was 2.0 on the medium containing IAA 1.0 mg l^{-1} + NAA 0.5 mg l^{-1} . However, it was decreased to 1.9 with the same concentration of NAA when the combination was with 1.0 mg l^{-1} IBA. Maximum 4.3 numbers of roots per shoot in seedling explants of guava on medium containing IBA 4.90 μM combined with activated charcoal 100 mg l^{-1} . was achieved by [11]. The highest number of roots per shoot may due to the combination of activated charcoal as activated charcoal acts as adsorbents to toxins, excess phytohormones and other phenolics.

Conclusions

It can be concluded from results that combination of BA 1.0 mg l^{-1} with 0.5 mg l^{-1} zeatin proved to be the best one in case of *in vitro* regeneration from seedlings explants of cv. safed (48 % shoot organogenesis). On the same medium maximum of 3.6 shoots

per explant was produced followed by BA 1.0 mg l^{-1} that produced 3.3 shoots per explant. In case of root organogenesis the best medium was IAA 1.5 mg l^{-1} combined with 1.0 mg l^{-1} NAA which induced root organogenesis in (76.6 %) micro shoots in 42-days. On the average, it gave 2.5 roots per explants.

Authors' contributions

Conceived and designed the experiments: A Rab, Performed the experiments: R Zamir, M Sajid & I Ahmad, Analyzed the data: R Zamir, A Rab & I Ahmad, Contributed reagents/materials/analysis tools: A Rab & I Ahmad, Wrote the paper: R Zamir, M Sajid & I Ahmad

References

1. Khushak A, Memon MA & Lashari MI (2009). Factors affecting guava production in Pakistan. *J Agric Res* 47(2): 201-209.
2. Amin MN & Jaiswal VS (1988). Micropropagation as an aid to rapid cloning of a guava cultivar. *Sci Hort* 36: 89-95.
3. Rai MK, Asthana P, Jaiswal VS & Jaiswal U (2010). Biotechnological advances in guava (*Psidium guajava* L.): Recent developments and prospects for future research. *Trees* 24: 1-12.

4. Zamir R, Khattak GSS, Mohammad T, Shah SA, Khan AJ & Ali N (2003). *In vitro* mutagenesis in guava (*Psidium guajava* L.). *Pak J Bot* 35(5):825-828.
5. Martin FW, Campbell CW & Roberte RM (1987). Perennial edible fruits of the tropics: An inventory, U.S. *Deptt Agri Res Ser Agr Hand book* 642.
6. Amin MN & Jaiswal VS (1993). *In vitro* response of apical bud explants from mature trees of jack fruit (*Artocarpus heterophyllus*). *Plant cell tiss & organ cult* 33: 59-65.
7. Pence VC & Sandoval JA (2002). Controlling contamination during *in vitro* collecting. A review of methods and chemicals used for controlling contamination in collected tissues. *IPGRI tech bulletin* 7: 30-40.
8. Loh CS & Rao AN (1989). Clonal propagation of guava (*Psidium guajava* L.) from seedling and grafted plants and adventitious shoot formation *in vitro*. *Scie Hort* 39: 31-39.
9. Papadatou P, Pontikis C, Eptimiadou E & Lydaki M (1990). Rapid multiplication of guava seedlings by *in vitro* shoot culture. *Scie Hort* 45: 99-103.
10. Yasseen M, Barringer SA, Schnell RJ & Splittoesser WE (1995). *In vitro* shoot proliferation of guava (*Psidium guajava* L.) from germinated seedlings. *Plant Cell Rep* 14: 525-528.
11. Goyal Y & Arya HC (1985). Tissue culture of desert tree- II. Clonal multiplication of *Ziziphus in vitro*. *J plant physiol* 199: 399-404.
12. Murashige T & Skoog FA (1962). Revised medium for rapid growth and bioassays with tobacco tissue cultures. *Phys Plant* 15: 473-497.
13. Chandra R, Mishra M & Bajpai A (2005). Biotechnological interventions for improvement of guava (*Psidium guajava* L.). Proceeding of 1st Int guava symp, CISH, Lucknow, India. pp. 19-25.
14. Mishra AK (2005). Important diseases of guava in India with special reference to wilt. In: Proc of 1st international guava symposium, CISH, Lucknow, India. pp.75-90.
15. Rai MK, Jaiswal VS & Jaiswal U (2009). Shoot multiplication and plant regeneration of guava (*Psidium guajava* L.) from nodal explants of *in vitro* raised plantlets. *J fruit and ornamental plant Res* 17(1): 29-38.
16. Ali N, Mulwa RMS, Norton MA & Skirvin RM (2003). Micropropagation of guava (*Psidium guajava* L.). *J Hort Sci & Biotech* 78: 739-741.
17. Kouidar M, Korban SS, Skirvin RM & Chu MC (1984). Influence of embryogenic dominance and polarity on adventitious shoot formation from apple cotyledons *in vitro*. *J Amer Soc Hort Sci* 109: 381-385.
18. Shahsavari E (2011). Impact of tryptophan and glutamin on the tissue culture of upland rice. *Plant soil envir.* 57(1): 7-10.
19. Jaiswal VS & Amin MN (1987). *In vitro* propagation of guava from shoot cultures of mature tree. *J Plant Physiol* 130: 7-12.
20. Haroun SA, Shukry WM & El-Sawy O (2010). Effect of asparagine or glutamin growth and metabolic changes in *phaseolus vulgaris* under *in vitro* conditions. *Biosci Res.* 7(1): 01-21.
21. Chandra R, Bajpai A, Gupta S & Tiwari RK (2004). Embryogenesis and plant regeneration from mesocarp of *Psidium guajava* L. *Indian J Biotech* 3(531): 246-248.
22. Das DK, Prakash NS & Balla-Sarin N (1999). Multiple shoot induction and plant regeneration in litchi (*litchi chinensis* Son). *Plant cell Rep* 18: 691-665.
23. Fuenmayor MED & Montero NJM (1997). *In vitro* clonal propagation of

- guava (*Psidium guajava* L.) from stem shoots of cv. Mara-7. *Acta Hort* 452: 47–51.
24. Malik SK, Chaudhury R & Kalia RK (2005). Rapid *in vitro* multiplication and conservation of *Garcinia indica*: A tropical medicinal tree species. *Sci Hort* 106: 539-553.
 25. Meghwal PR, Singh SK & Sharma HC (2003). Micropropagation of Aneuploid guava. *Indian J Hort* 60: 29-33.
 26. Mishra DS, Tiwari JP & Lal S (2005). *In vitro* cloning of guava (*Psidium guajava* L.) cv. Pant Prabhat. In: Proc of 1st International guava symposium, CISH, Lucknow, India; pp. 32-33.
 27. Singh SK, Meghwal PR, Sharma HC & Singh SP (2002). Direct shoot organogenesis on explants from germinated seedlings of *Psidium guajava* L. cv. Allahabad Safeda. *Sci Hort* 95: 213-221.
 28. Amin MN & Jaiswal VS (1987). Rapid clonal propagation of guava through *in vitro* shoot proliferation on nodal explants of mature trees. *Plant Cell Tiss & Organ Cult* 9: 235-243.
 29. Roy SK, Islam MS, Sen J & Hadiuzzaman S (1992). Effects of auxins, sucrose and agar on *in vitro* rooting of callus induced micro shoots of jackfruit (*Artocarpus heterophyllus*). *Bangladesh J Bot* 21: 93:98.
 30. Amin MN, Razzaque MA & Akhtar S (1992). Axillary shoot proliferation and adventitious rooting *in vitro* of response of carambola (*Averrhoa carambola* L.). *Plant Tiss Cult* 2: 7-13.
 31. Amin MN, Islam MN & Azad AK (1999). Regeneration of plantlets *in vitro* from the seedling explants of pomegranate (*Punica granatum*). *Plant Tiss Cult* 9(1): 53-61.
 32. MacRae S, Van J & Den (1990). *In vitro* culture of *Eucalyptus grandis*: Effect of gelling agents on propagation. *J Plant Physiol* 137: 249-251.
 33. Singh D, Mohammed S & Shukla AK (2005). Micropropagation of guava as influenced by season, source and pretreatment to explant. In: Proc of 1st international guava symposium, CISH, Lucknow, Ind pp. 33.